

SURVEY DATES

SURVEY LOCATIONS

 07/17/2006 thru 07/28/2006 BUCHANAN, DICKENSON, RUSSELL AND TAZEWEEL
 COUNTIES, VIRGINIA

DEMOGRAPHICS

Estimated current employment:	1,754	Estimated 5-year employment:	1,799
Number of miners examined:	305	5-YEAR miners examined:	611
Survey participation rate:	17%	5-year participation rate:	34%

	N	Mean	Range
Tenure:	299	25	0 - 42
Age:	305	48	22 - 60
Tenure at the Face:	284	22	1 - 42

PREVALENCE OF DISEASE: n=305

Pneumoconiosis:	27 (9%)	PMF:	6 (2%)	Advanced:	10 (3%)
-----------------	----------	------	---------	-----------	----------

EXPECTED PREVALENCE OF DISEASE FOR WORKERS WITH 15 OR MORE TENURE
 YEARS AND THE AVERAGE LEVEL OF DUST FOR THE COUNTIES OF 0.881 MG/M3: n=255

Pneumoconiosis:	12 (5%)	PMF:	2 (1%)	Advanced:	4 (1%)
-----------------	----------	------	---------	-----------	---------

DISTRIBUTION OF RADIOGRAPHIC FINDINGS: n=305

Profusion of Small Opacities	Large Opacities	Symbols
0/-: 0 (0%)	O: 299 (98%)	ax: 10 (3%)
0/0: 253 (83%)	A: 2 (1%)	
0/1: 25 (8%)	B: 4 (1%)	
1/0: 3 (1%)	C: 0 (0%)	
1/1: 7 (2%)		
1/2: 8 (3%)		
2/1: 2 (1%)		
2/2: 4 (1%)		
2/3: 2 (1%)		
3/2: 0 (0%)		
3/3: 1 (0%)		
3/+: 0 (0%)		

PREVALENCE OF CLINICAL FINDINGS: n=305

Chronic cough:	78 (26%)	Chest tightness:	106 (35%)
Phlegm:	92 (30%)	Chronic bronchitis:	23 (8%)
Wheezing:	88 (29%)	Emphysema:	9 (3%)
Attacks of wheezing:	127 (42%)	Pneumonia:	50 (16%)
Dyspnea (on level):	131 (43%)	Asthma:	19 (6%)

Dyspnea (up hill):	191 (63%)	Tuberculosis:	0 (0%)
--------------------	------------	---------------	---------

Report Date: 06/21/2007

Page 2

Smoking status: n=298

Never		Former		Current	
Smoked:	140 (47%)	Smoker:	89 (30%)	Smoker:	69 (23%)

SPIROMETRY RESULTS: n=301

Obstructive:	22 (7%)	Restrictive:	9 (3%)	Mixed:	3 (1%)
--------------	----------	--------------	---------	--------	---------

METHODS

DEMOGRAPHICS

SURVEY PARTICIPATION RATE: count of the number of miners with a valid x-ray during the survey period (NUMBER OF MINERS EXAMINED) divided by the count of the names on the most recent mine rosters submitted to NIOSH by the mine operators (ESTIMATED CURRENT EMPLOYMENT).

5-YEAR PARTICIPATION RATE: count of the number miners with a valid x-ray during the 5-year period preceding the date of the end of the survey (5-YEAR MINERS EXAMINED) divided by the count of the names on the mine rosters that accompanied the mine plans that were submitted to NIOSH by the miner operators during the 5-year period preceding the date of the end of the survey (ESTIMATED 5-YEAR EMPLOYMENT).

TENURE - MEAN: summation of the reported years in various underground mining occupations divided by count of participating miners with a completed work history (N).

AGE - MEAN: summation of the calculated age at time of the x-ray divided by count of participating miners who provided their date of birth (N).

TENURE AT THE FACE - MEAN: summation of the reported years of working at the face divided by count of participating miners who reported working at the face (N).

PREVALENCE OF DISEASE

PNEUMOCONIOSIS: number of x-rays with a final determination (consensus based upon NIOSH procedures) of the International Labour Office (ILO) small opacity profusion category 1/0 or higher or with the presence of large opacities, divided by the number of participating miners with a valid x-ray.

PMF: number of x-rays with a final determination of ILO large opacity category "A", "B", or "C" divided by the number of participating miners with a valid x-ray.

ADVANCED: number of x-rays with a final determination of ILO small opacity profusion category 2/1 or higher and presence of symbol "ax" (coalescence of small opacities) or ILO large opacity category of "A", "B", or "C" divided by the number of participating miners with a valid x-ray.

EXPECTED PREVALENCE

PNEUMOCONIOSIS, PMF & ADVANCED: count and percent of expected cases of pneumoconiosis, PMF, and advanced disease for a given mean respirable dust level (concentration) and for the specific type of coal in the region, calculated according to Attfield MD, Morring K. Am Ind Hyg Assoc J 1992; 53:486-492.

METHODS (continued)

DISTRIBUTION OF RADIOGRAPHIC FINDINGS

PROFUSION OF SMALL OPACITIES, LARGE OPACITIES & SYMBOLS: count and percent of the x-rays using the final determination of the ILO small opacity, the final determination of the ILO large opacity, and symbol "ax".

PREVALENCE OF CLINICAL FINDINGS

SELF-REPORTED: count and percent of self-reported symptoms and/or medical conditions based on the participating miners who completed the health questionnaire survey.

SMOKING STATUS

NEVER SMOKER, FORMER SMOKER, & CURRENT SMOKER: count and percent of participating miners who completed the health questionnaire survey miners reporting each smoking history.

SPIROMETRY RESULTS

OBSTRUCTIVE, RESTRICTIVE, & MIXED: number and percent of miners who participated in the spirometry testing and whose results showed one of the following categories of respiratory impairment using forced expiratory volume in one second (FEV1) and forced vital capacity (FVC) ratio:

OBSTRUCTIVE - $FEV1/FVC < \text{lower limit of normal (LLN)}$ and $FVC \geq LLN$
RESTRICTIVE - $FEV1/FVC > LLN$ and $FVC < LLN$
MIXED - $FEV1/FVC < LLN$ and $FVC < LLN$